

UNITED STATES PATENT APPLICATION

For

ALLANTOIN-CONTAINING SKIN CREAM

Assignee: Alwyn Company, Inc.
Inventor: Elliott Farber

OPPENHEIMER

OPPENHEIMER WOLFF & DONNELLY LLP

2029 Century Park East, Suite 3800
Los Angeles, California 90067
Tel (310) 788-5000
Fax (310) 788-5100

Attorney Matter No. 14358-315

ALLANTOIN-CONTAINING SKIN CREAM

CROSS REFERENCES

[0001] This application is a continuation-in-part application of Application Serial No. 09/758,781, entitled "Allantoin-Containing Skin Cream," filed January 11, 2001 by Elliott Farber, which in turn was a continuation-in-part application of 09/570,266, entitled "Allantoin-Containing Skin Cream," filed May 12, 2000 by Elliott Farber, which in turn is a continuation-in-part application of Application Serial No. 09/360,095, entitled "Oil-in-Water Emulsion With Improved Stability," filed July 23, 1999 by Elliott Farber, now U.S. Patent No. 6,281,236, issued August 28, 2001. All three of these prior applications are hereby incorporated in their entirety by this reference.

BACKGROUND OF THE INVENTION

[0002] *Field of the Invention:* This invention is directed to a skin cream containing allantoin in an oil-in-water emulsion with improved stability.

[0003] *General Background and State of the Art:* Allantoin is a commonly used ingredient in cosmetic applications, particularly for skin creams, where it exerts a skin protective function. Many such cosmetic compositions and other compositions are prepared as emulsions, particularly oil-in-water emulsions. One emulsifier system used with such compositions is a combination of sodium lauryl sulfate and beeswax. Although solutions of sodium lauryl sulfate are alkaline with an approximate pH of 9.5, the simultaneous use of beeswax with its organic acids produces a complex and neutralized system with a pH of about 6.8 to about 7.5. However, in such a system with a pH range of 6.8 to 7.5, allantoin degrades significantly with time and in accelerated stability tests at 40°C. Because cosmetics and other preparations designed for application to the skin are typically stored by users at room temperature, and room temperatures can fluctuate with climatic conditions, such a degree of stability is undesirable. Therefore, there is a need for an oil-in-water emulsified composition containing allantoin in which the stability of allantoin is increased.

[0004] In particular, there is a need for compositions that are suitable for treating a number of severe and difficult-to-treat skin conditions. One of these skin conditions is

epidermolysis bullosa. This is a severe genetic skin disorder in which the skin breaks down and large blisters appear. These blisters are difficult to treat by conventional means. Other skin diseases for which improved treatments are needed are pressure ulcers, decubitus ulcers or bed sores, and diabetic ulcers, and milia, as well as other conditions affecting the skin and having an inflammatory component such as eczema, urticaria, atopic dermatitis, contact dermatitis, arthritis, gout, and lupus erythematosus. Therefore, improved compositions that are suitable for treating these diseases are needed.

INVENTION SUMMARY

[0005] In general, a composition according to the present invention comprises an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) at least one anionic or nonionic emulsifier that is substantially hydrophilic and is soluble in water.

[0006] The pH of the composition is in a range of from about 3.0 to about 6.0. This pH range stabilizes the allantoin and makes storage of the composition for extended periods practical. Preferably, the pH of the composition is in a range of from about 5.0 to about 6.0.

[0007] Typically, the at least one anionic or nonionic emulsifier is selected from the group consisting of:

- (1) an acidic anionic polymer;
- (2) an anionic emulsifier selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate;
- (3) a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose chain length ranges from 8 to 22 carbon atoms;
- (4) glyceryl stearate;
- (5) cetyl alcohol;
- (6) stearic acid;
- (7) sodium stearyl lactylate;
- (8) sodium isostearyl lactylate;

(9) triethanolamine stearate;

(10) a polyethylene glycol ether of cetearyl alcohol wherein the number of polyethylene glycol moieties in the ether is from 6 to 40; and

(11) an acidic wax.

[0008] The composition can further comprise a carbohydrate polymer selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Typically, the carbohydrate polymer is galactoarabinan.

[0009] Acidic waxes are those waxes having acidic groups that can be neutralized with alkaline materials such as hydroxides, alkoxides, unprotonated amines, and/or salts of strong bases and weak acids, such as sodium acetate. Upon neutralization, such waxes can act as emulsifiers or coemulsifiers. Particularly preferred acidic waxes include beeswax, carnauba wax, candelilla wax, siliconyl beeswax, siliconyl carnauba wax, and synthetic acidic waxes. A particularly preferred acidic wax is beeswax.

[0010] One embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) an acidic anionic polymer; and
 - (b) a polyethylene glycol ester of stearic acid.

[0011] The pH of the composition is from about 3.0 to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 6.0.

[0012] Typically, in this embodiment, the acidic anionic polymer is a carboxypolymethylene polymer.

[0013] The composition can further comprise a carbohydrate polymer. Typically, the carbohydrate polymer is selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Preferably, the carbohydrate polymer is galactoarabinan.

[0014] Another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and

(2) an emulsifier system comprising:

(a) an acidic anionic polymer; and

(b) an anionic emulsifier that is substantially hydrophilic and is soluble in water, the pH of the composition being adjusted to a range of from about 3.0 to about 6.0. Preferably, the pH of the composition is adjusted to a range of from about 5.0 to about 6.0.

[0015] The anionic emulsifier can be selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate. Preferably, the anionic emulsifier is sodium lauryl sulfate.

[0016] The composition can further comprise a carbohydrate polymer selected from the group consisting of galactoarabinan, polygalactose and polyarabinose. Preferably, the carbohydrate polymer is galactoarabinan.

[0017] Another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin;

(2) an emulsifier system comprising an acidic anionic polymer; and

(3) a base to adjust the pH of the composition to a value in a range from about 3.0 to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 5.5.

[0018] Preferably, the acidic anionic polymer is a carboxypolymethylene polymer. Preferably, the base is an organic base such as triethanolamine.

[0019] In another embodiment, the composition comprises an oil-in-water emulsion comprising:

(1) allantoin;

(2) an emulsifier system comprising:

(a) an acidic anionic polymer; and

(b) a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms, wherein the pH of the composition is from about 3.0 to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 6.0.

[0020] In this embodiment, the emulsifier system can further comprise glyceryl stearate.

[0021] This embodiment also can include a carbohydrate polymer as described above.

[0022] In yet another embodiment of the present invention, a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin;
- (2) an emulsifier system comprising:
 - (a) sodium stearyl lactylate;
 - (b) sodium isostearyl lactylate;
 - (c) optionally, triethanolamine stearate;
 - (d) optionally, at least one nonionic emulsifier selected from the group

consisting of a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms; and

- (3) an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from 5.0 to about 5.8.

[0023] Typically, the acid is citric acid.

[0024] Still another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising at least one polyethyleneglycol ether of cetearyl alcohol, wherein the number of polyethylene glycol moieties in the polyethyleneglycol ether of cetearyl alcohol is from 6 to 40; and
- (3) an acid to adjust the pH of the composition to a range of from about 5.0 to about 5.8.

[0025] For this embodiment, the acid is typically also citric acid.

[0026] For this embodiment, the emulsifier system typically comprises cetareth-25 and cetareth-6.

[0027] Yet another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin;
- (2) an emulsifier system comprising:
 - (a) a polyethylene glycol ester of stearic acid; and
 - (b) glyceryl stearate; and

(3) an acid to adjust the pH of the composition to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8.

[0028] Typically, in this embodiment, the number of ethylene glycol moieties in the polyethylene glycol ester of stearic acid is from 25 to 100. Preferably, the polyethylene glycol ester of stearic acid is PEG-100 stearate. Typically, in this embodiment, the acid is citric acid.

[0029] Still another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin;
- (2) a carbohydrate polymer; and
- (3) an emulsifier system comprising:
 - (a) an acidic wax; and
 - (b) an anionic emulsifier that is substantially hydrophilic and is soluble in

water;

[0030] The pH of the composition is in a range from about 3.0 to about 6.0. Preferably, the pH is in the range of from about 5.0 to about 6.0.

[0031] The carbohydrate polymer in this embodiment is as described above.

[0032] Typically, the anionic emulsifier that is substantially hydrophilic and soluble in water is selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate.

Preferably, the anionic emulsifier is sodium lauryl sulfate.

[0033] The acidic wax in this embodiment is as described above. Typically, the acidic wax is beeswax, carnauba wax, candelilla wax, siliconyl beeswax, siliconyl carnauba wax, or a synthetic acidic wax. Preferably, the acidic wax is beeswax.

[0034] The composition can further comprise citric acid to adjust the pH.

[0035] Still another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin in a concentration of at least about 2.5% and;

(2) an emulsifier system comprising:

- (a) an acidic wax; and
- (b) an anionic emulsifier that is substantially hydrophilic and is soluble in

water.

[0036] The pH of the composition is in a range from about 3.0 to about 6.0.

[0037] Typically, the anionic emulsifier that is substantially hydrophilic and soluble in water is selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate.

Preferably, the anionic emulsifier is sodium lauryl sulfate.

[0038] The acidic wax in this embodiment is as described above. Typically, the acidic wax is beeswax, carnauba wax, candelilla wax, siliconyl beeswax, siliconyl carnauba wax, or a synthetic acidic wax. Preferably, the acidic wax is beeswax.

[0039] The composition can further comprise citric acid to adjust the pH.

[0040] For all of these embodiments, the composition can comprise one or more additional ingredients as described below.

[0041] The composition can comprise an emollient component comprising at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

[0042] The composition can comprise an emollient such as butylated hydroxytoluene.

[0043] The composition can comprise herbal extracts such as one or more of St. John's wort extract, witch hazel extract, chamomile extract, and arnica extract.

[0044] The composition can comprise a preservative component comprising at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea.

[0045] The composition can comprise a chelating agent such as tetrasodium EDTA.

[0046] The composition can comprise a solvent component comprising at least one solvent selected from the group consisting of propylene glycol, butylene glycol and glycerin. Preferably, the solvent component is propylene glycol.

[0047] Ranges of compositions are disclosed for each of these embodiments.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0048] A composition according to the present invention comprises an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) at least one anionic or nonionic emulsifier that is substantially hydrophilic and is soluble in water.

[0049] The composition can further include other ingredients, such as a chelating agent to bind metal ions that might accelerate degradation of the composition. A particularly preferred chelating agent is EDTA. The EDTA can be added in various acid or salt forms depending on the pH of the composition, such as EDTA itself, disodium EDTA, or tetrasodium EDTA.

[0050] The pH of the composition is in a range of from about 3.0 to about 6.0. This pH range stabilizes the allantoin and makes storage of the composition for extended periods practical. Preferably, the pH is in a range of from about 5.0 to about 6.0. In some cases, depending on the ingredients used in the composition, narrower pH ranges, such as from about 5.0 to about 5.8, or from about 5.0 to about 5.5, are desirable.

[0051] Typically, the at least one anionic or nonionic emulsifier is selected from the group consisting of:

- (1) an acidic anionic polymer;
- (2) an anionic emulsifier selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate;
- (3) a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose chain length ranges from 8 to 22 carbon atoms;

- (4) glyceryl stearate;
- (5) cetyl alcohol;
- (6) stearic acid;
- (7) sodium stearyl lactylate;
- (8) sodium isostearyl lactylate;
- (9) triethanolamine stearate;
- (10) a polyethylene glycol ether of cetearyl alcohol wherein the number of polyethylene glycol moieties in the ether is from 6 to 40; and
- (11) an acidic wax.

[0052] The composition can further comprise a carbohydrate polymer selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Typically, the carbohydrate polymer is galactoarabinan.

[0053] Acidic waxes are those waxes having acidic groups that can be neutralized with alkaline materials such as hydroxides, alkoxides, unprotonated amines, and/or salts of strong bases and weak acids, such as sodium acetate. Upon neutralization, such waxes can act as emulsifiers or coemulsifiers. Particularly preferred acidic waxes include beeswax, carnauba wax, candelilla wax, siliconyl beeswax, siliconyl carnauba wax, and synthetic acidic waxes. Examples of synthetic acidic waxes are syncrowaxes marketed by Croda, Inc.

[0054] One embodiment of an improved composition containing allantoin is a composition that comprises an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) an acidic anionic polymer; and
 - (b) a polyethylene glycol ester of stearic acid.

[0055] The pH of the composition is adjusted to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 6.0. The pH is adjusted with sodium hydroxide or another base as required.

[0056] The acidic anionic polymer is preferably a carboxypolymethylene polymer. Such polymers are marketed under the brand names "Carbomer" and "Carbopol." A suitable

carboxypolymethylene polymer is marketed by B.F. Goodrich under the brand name "Carbomer." This is a slightly cross-linked polyacrylic acid that is from 1% to 2% cross-linked by allylsucrose or allylpentaerythritol with the polyacrylic acid. The resulting molecular weight range of this polymer is from about 2×10^6 daltons to about 1×10^9 daltons. The average molecular weight of this polymer is about 4×10^6 daltons.

[0057] Preferably, the concentration of the carboxypolymethylene polymer is from about 0.5 percent to about 2 percent of the composition.

[0058] The composition can further comprise a carbohydrate polymer. Preferably, the carbohydrate polymer is galactoarabinan. Galactoarabinan is derived from trees of the genus Larix (larch) and is a hemicellulosic product easily extractable by water in a pure form. Galactoarabinan has been consumed by humans in common foods such as carrots, tomatoes, maple syrup, soybeans, and wheat flour, among others. The molecular weight of the galactoarabinan is about 20,000. A suitable source of galactoarabinan is Larex, Inc. (White Bear Lake, MN). Typically, the composition contains from about 1 percent to about 25 percent of galactoarabinan. Preferably, the composition contains from about 2 percent to about 10 percent of the carbohydrate polymer.

[0059] This embodiment of a composition according to the present invention can further include other ingredients. For example, the composition can include an emollient component for smoothness. The emollient component can include at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

[0060] The composition can also include an antioxidant to prevent rancidity of ingredients such as cod liver oil. A preferred antioxidant is butylated hydroxytoluene (BHT). Other antioxidants such as butylated hydroxyanisole (BHA) can be used, alternatively or in addition to BHT.

[0061] The composition can further include a solvent component. Typically, the solvent component can include at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0062] The composition can further include a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the preservative component comprises methylparaben, propylparaben, and diazolidinyl urea. Other preservatives can also be used.

[0063] The composition can further include fragrance. The use of fragrance is well known in the cosmetic art and in the art of over-the-counter drug formulation, and many suitable fragrances are known in the art. The stability and function of the cream is not altered by the presence or absence of fragrance.

[0064] Optionally, this embodiment of the composition can further include herbal extracts. The herbal extracts can include one or more of St. John's wort extract, witch hazel extract, chamomile extract, and arnica extract. However, these herbal extracts are typically omitted in this embodiment.

[0065] This embodiment of the composition can optionally further include other components, such as proteins, humectants, other preservatives, essential oils, other vitamins, colorants, hydroxyacids, other plant extracts, chelators, sunscreens, sodium hyaluronate, lipids, fatty acids, thickeners, panthenol, and the like. The use of such components is conventional in the cosmetic art and in the over-the-counter drug art. Typical sunscreens are octyl methoxycinnamate and benzophenone-3.

[0066] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention with a pH range of from about 5.0 to about 6.0. For this and other ranges, preferred concentrations, and optimum concentrations of specific ingredients for other embodiments as given below, all percentages are weight percentages unless otherwise specified.

[0067] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 85.0% of this embodiment of the composition. In one alternative, in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 69.95% of the composition. In another alternative, in which the optimum concentration of

allantoin is about 9.00% of the composition, the optimum concentration of water is about 62.45% of the composition.

[0068] The carboxypolymethylene polymer can comprise from about 0.30% to about 3.0% of this embodiment of the composition. Preferably, the carboxypolymethylene polymer comprises from about 0.50% to about 2.0% of this embodiment of the composition. An optimum concentration of the carboxypolymethylene polymer is about 0.85% of this embodiment of the composition.

[0069] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0070] PEG-100 stearate can comprise from about 0.25% to about 2.5% of this embodiment of the composition. Preferably, PEG-100 stearate comprises from about 0.50% to about 2.0% of this embodiment of the composition. An optimum concentration of PEG-100 stearate is about 1.50% of this embodiment of the composition.

[0071] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0072] Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. A preferred concentration of cetyl alcohol is from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.20% of this embodiment of the composition.

[0073] Stearyl alcohol can comprise from about 0.5% to about 6.0% of this embodiment of the composition. A preferred concentration of stearyl alcohol is from about 0.75% to about 5.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 1.50% of this embodiment of the composition.

[0074] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this

embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

[0075] Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0076] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. A preferred concentration of methylparaben is from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0077] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0078] Diazolidinyl urea can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.15% of this embodiment of the composition.

[0079] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of this embodiment of the composition.

[0080] Fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of the composition. As indicated above, fragrance can be omitted, and it may be desirable to

omit fragrance in circumstances in which the composition is intended for use on sensitive individuals or individuals who may undergo an allergic reaction to fragrance.

[0081] Triethanolamine can comprise from about 0.05% to about 3.0% of this embodiment of the composition to adjust the pH. A preferred concentration of triethanolamine is from about 0.20% to about 2.0% of this embodiment of the composition. An optimum concentration of triethanolamine is about 0.80% of this embodiment of the composition.

[0082] In another alternative embodiment of the composition, the emulsifier can be an anionic emulsifier that is substantially hydrophilic and is soluble in water. In this embodiment, the anionic emulsifier replaces the polyethylene glycol ester of stearic acid. This embodiment further includes the acidic anionic polymer such as carboxypolymethylene. Optionally, but preferably, this alternative embodiment of the composition includes the carbohydrate polymer such as galactoarabinan.

[0083] The anionic emulsifier that is substantially hydrophilic and soluble in water can be selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate. A particularly preferred anionic emulsifier is sodium lauryl sulfate.

[0084] Commercially available preparations of sodium lauryl sulfate contain sufficient excess sodium hydroxide so that they have a pH of about 10.0. This sodium hydroxide can be used to adjust the pH when the anionic emulsifier is sodium lauryl sulfate; in this alternative, no additional alkali may be needed. When another anionic emulsifier is used, additional alkali may be required to adjust the pH.

[0085] In yet another alternative embodiment of the composition, the emulsifier system comprises the acidic anionic polymer as described above and a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms.

[0086] Preferably, the acidic anionic polymer is carboxypolymethylene as described above.

[0087] This alternative embodiment of the composition can further include glyceryl stearate in the emulsifier system.

[0088] This embodiment of the composition has a pH from about 3.0 to 6.0, adjusted as necessary, typically with an acid. The acid can be an organic acid, an inorganic acid, or a mixture of both. Preferably, the pH is from about 5.0 to about 6.0.

[0089] This embodiment of the composition can further comprise a carbohydrate polymer such as galactoarabinan as described above.

[0090] In this embodiment of the composition, preferred organic acids include organic acids whose carbon chain length ranges from 2 to 22 carbon atoms and that can be monocarboxylic, dicarboxylic, or tricarboxylic acids. The acids can be aliphatic or aromatic. Particularly preferred organic acids include citric acid, ascorbic acid, glycolic acid, lactic acid, benzoic acid, and salicylic acid. A most particularly preferred organic acid is citric acid.

[0091] Typically, in this embodiment of the composition, the inorganic acid is a strong acid. It can be a monoprotic, diprotic, or triprotic acid. Particularly preferred inorganic acids include hydrochloric acid, sulfuric acid, and phosphoric acid.

[0092] This alternative embodiment of the composition can further include other ingredients as described above, including an emollient component, an antioxidant, a solvent component, a chelating agent, herbal extracts, a preservative, and fragrance.

[0093] The composition can further include other components, such as proteins, humectants, other preservatives, essential oils, other vitamins, colorants, hydroxyacids, other plant extracts, sunscreens, sodium hyaluronate, lipids, fatty acids, thickeners, panthenol, and the like. The use of such components is conventional in the cosmetic art and in the over-the-counter drug art. Typical sunscreens are octyl methoxycinnamate and benzophenone-3.

[0094] In yet another embodiment of the composition, the emulsifier system comprises the acidic anionic polymer described above; one example of this acidic anionic polymer is marketed as Carbomer. In this embodiment, the pH is adjusted with an organic or inorganic base to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.5. A preferred organic base is triethanolamine. A preferred inorganic base is sodium hydroxide. In general, it is preferred to use an organic base such as triethanolamine.

[0095] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0096] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one solvent selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

[0097] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole can also be used.

[0098] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben. As indicated above, other preservatives can also be used.

[0099] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0100] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention where the pH is from about 5.0 to about 5.5.

[0101] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 80.0% of this embodiment of the composition. In one alternative, in which the optimum concentration of allantoin is about 1.50% of the composition, an optimum concentration of water is about 73.55% of this embodiment of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 66.05% of the composition.

[0102] The carboxypolymethylene polymer can comprise from about 0.40% to about 3.0% of this embodiment of the composition. Preferably, the carboxypolymethylene polymer comprises from about 0.5% to about 2.0% of this embodiment of the composition. An optimum concentration of the carboxypolymethylene polymer is about 1.00% of the composition.

[0103] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, the propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of the propylene glycol is about 5.70% of this embodiment of the composition.

[0104] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.0% of this embodiment of the composition.

[0105] Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.00% of this embodiment of the composition.

[0106] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

[0107] Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.30% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0108] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0109] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0110] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of the composition.

[0111] Fragrance, if present, can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, if present, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance, if present, is about 0.20% of this embodiment of the composition.

[0112] Triethanolamine, as a 95% solution, can comprise from about 0.05% to about 3.0% of this embodiment of the composition to adjust the pH to a value in the range of from about 5.0 to about 5.5. Preferably, triethanolamine comprises from about 0.20% to about 2.0% of this embodiment of the composition to adjust the pH as indicated. An optimum concentration of triethanolamine is about 0.80% of the composition to adjust the pH as indicated.

[0113] Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) cetyl alcohol; and
 - (b) stearic acid.

[0114] In this embodiment, the pH is adjusted to a range of from about 3.0 to about 6.0 by addition of a quantity of a weak organic base. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The weak organic base can be an amine-containing base such as ethanolamine, diethanolamine, or triethanolamine. A preferred organic base is triethanolamine.

[0115] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0116] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one solvent selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

[0117] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole can also be used.

[0118] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben. As indicated above, other preservatives can alternatively be used.

[0119] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream is not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0120] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is in the range of from about 5.0 to about 5.8.

[0121] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 85.0% of this embodiment of the composition. In one alternative, in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 71.70% of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 64.20% of the composition.

[0122] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0123] Triethanolamine can comprise from about 0.2% to about 4.0% of this embodiment of the composition. Preferably, triethanolamine comprises from about 0.5% to about 3.0% of this embodiment of the composition. An optimum concentration of triethanolamine is about 1.25% of this embodiment of the composition.

[0124] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0125] Cetyl alcohol can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 6.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.50% of this embodiment of the composition.

[0126] Stearic acid can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, stearic acid comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of stearic acid is about 2.50% of this embodiment of the composition.

[0127] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.50% to about 5.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.0% of this embodiment of the composition.

[0128] Butylated hydroxytoluene can comprise from about 0.1% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.2% to about 0.8% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0129] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0130] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0131] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of the composition.

[0132] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0133] Still another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) sodium stearoyl lactylate;
 - (b) sodium isostearoyl lactylate;
 - (c) optionally, triethanolamine stearate;
 - (d) optionally, at least one nonionic emulsifier selected from the group

consisting of a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms.

[0134] Sodium stearoyl lactylate is the sodium salt of the stearic acid ester of lactyl lactate. Sodium isostearoyl lactylate is the sodium salt of the isostearic acid ester of lactyl lactate.

[0135] In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

[0136] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0137] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

[0138] This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

[0139] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben. As indicated above, other preservatives can be used.

[0140] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole can also be used.

[0141] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0142] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is in the range of from about 5.0 to about 5.8.

[0143] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 80.0% of this embodiment of the composition. In one alternative, in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 73.72% of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 66.22% of the composition.

[0144] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0145] Citric acid can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of citric acid is about 0.18% of this embodiment of the composition.

[0146] Sodium stearyl lactylate can comprise from about 0.30% to about 3.0% of this embodiment of the composition. Preferably, sodium stearyl lactylate comprises from about 0.50% to about 2.50% of this embodiment of the composition. An optimum concentration of sodium stearyl lactylate is about 1.00% of this embodiment of the composition.

[0147] Sodium isostearyl lactylate can comprise from about 0.05% to about 1.0% of this embodiment of the composition. Preferably, sodium isostearyl lactylate comprises from about 0.10% to about 0.70% of this embodiment of the composition. An optimum concentration of sodium isostearyl lactylate is about 0.25% of this embodiment of the composition.

[0148] Tetrasodium EDTA can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

[0149] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0150] Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.80% of this embodiment of the composition.

[0151] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.0% of this embodiment of the composition.

[0152] Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0153] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0154] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0155] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of the composition.

[0156] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0157] Still another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising at least one polyethyleneglycol ether of cetearyl alcohol.

[0158] In polyethylene glycol ethers of cetearyl alcohol suitable for use in compositions according to the present invention, the number of ethylene glycol moieties can range from 6 to 40, e.g., $R(OCH_2CH_2)_{25}OH$ where R is $CH_3(CH_2)_{16-18}$. In one preferred embodiment of the present invention, the emulsifier system comprises both cetareth-25 and cetareth-6, i.e., polyethylene glycol ethers of cetearyl alcohol with 25 and 6 ethylene glycol units respectively.

[0159] In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

[0160] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0161] This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

[0162] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the

emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

[0163] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole can alternatively be used.

[0164] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the preservative component comprises all of methylparaben, propylparaben, and diazolidinyl urea. As indicated above, other preservatives can alternatively be used.

[0165] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0166] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is adjusted to a range of from about 5.0 to about 5.8.

[0167] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 55.0% to about 75.0% of this embodiment of the composition. In one alternative, in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 66.33% of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 58.83% of the composition.

[0168] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.2% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0169] Tetrasodium EDTA can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about

0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

[0170] Ceteareth-25 can comprise from about 0.50% to about 4.0% of this embodiment of the composition. Preferably, ceteareth-25 comprises from about 2.0% to about 3.5% of this embodiment of the composition. An optimum concentration of ceteareth-25 is about 2.60% of this embodiment of the composition.

[0171] Citric acid can comprise from about 0.04% to about 0.40% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of citric acid is about 0.12% of this embodiment of the composition.

[0172] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0173] Cetyl alcohol can comprise from about 3.0% to about 10.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 3.5% to about 7.5% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.30% of this embodiment of the composition.

[0174] Stearyl alcohol can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 2.0% to about 4.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 3.50% of this embodiment of the composition.

[0175] Ceteareth-6 can comprise from about 0.5% to about 4.0% of this embodiment of the composition. Preferably, ceteareth-6 comprises from about 1.0% to about 3.0% of this embodiment of the composition. An optimum concentration of ceteareth-6 is about 1.80% of this embodiment of the composition.

[0176] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

[0177] Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0178] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0179] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0180] Diazolidinyl urea can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.15% of this embodiment of the composition.

[0181] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of the composition.

[0182] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0183] Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:

- (a) a polyethylene glycol ester of stearic acid; and
- (b) glyceryl stearate.

[0184] Typically, the number of ethylene glycol moieties in the polyethylene glycol ester of stearic acid is from 25 to 100. Two preferred polyethylene glycol esters of stearic acid for use in this embodiment of compositions according to the present invention are PEG-40 stearate and PEG-100 stearate, with 40 and 100 ethylene glycol moieties respectively. A particularly preferred polyethylene glycol ester of stearic acid is PEG-100 stearate.

[0185] In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

[0186] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component is one or more of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0187] This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

[0188] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

[0189] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole can alternatively be used.

[0190] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the

preservative component comprises all of methylparaben, propylparaben, and diazolidinyl urea. As indicated above, other preservatives can alternatively be used.

[0191] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0192] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is in the range of from about 5.0 to about 5.8.

[0193] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 55.0% to about 80.0% of this embodiment of the composition. In one alternative in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 67.86% of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 60.36% of the composition.

[0194] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.30% to about 7.00% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0195] Tetrasodium EDTA can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

[0196] Citric acid can comprise from about 0.04% to about 0.40% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of citric acid is about 0.14% of this embodiment of the composition.

[0197] PEG-100 stearate can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, PEG-100 stearate comprises from about

1.50% to about 3.00% of this embodiment of the composition. An optimum concentration of PEG-100 stearate is about 2.60% of this embodiment of the composition.

[0198] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0199] Cetyl alcohol can comprise from about 2.0% to about 10.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.50% to about 7.50% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.00% of this embodiment of the composition.

[0200] Stearyl alcohol can comprise from about 1.0% to about 4.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 1.0% to about 3.5% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 2.50% of this embodiment of the composition.

[0201] Glyceryl stearate can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, glyceryl stearate comprises from about 2.0% to about 4.0% of this embodiment of the composition. An optimum concentration of glyceryl stearate is about 2.50% of this embodiment of the composition.

[0202] Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

[0203] Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0204] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to 0.40% of

this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0205] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0206] Diazolidinyl urea can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.20% of this embodiment of the composition.

[0207] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of this embodiment of the composition.

[0208] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0209] Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin;
- (2) a carbohydrate polymer; and
- (3) an emulsifier system comprising:
 - (a) an acidic wax; and
 - (b) an anionic emulsifier that is substantially hydrophilic and is soluble in

water.

[0210] Acidic waxes are those waxes having acidic groups that can be neutralized with alkaline materials such as hydroxides, alkoxides, unprotonated amines, and/or salts of

strong bases and weak acids, such as sodium acetate. Upon neutralization, such waxes can act as emulsifiers or coemulsifiers. Preferred acidic waxes include beeswax, carnauba wax, candelilla wax, siliconyl beeswax, siliconyl carnauba wax, and synthetic acidic waxes. Examples of synthetic acidic waxes are syncrowaxes marketed by Croda, Inc. A particularly preferred acid wax is beeswax.

[0211] The carbohydrate polymer is typically selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Preferably, the carbohydrate polymer is galactoarabinan.

[0212] The anionic emulsifier that is substantially hydrophilic and soluble in water can be selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate. A particularly preferred anionic emulsifier is sodium lauryl sulfate.

[0213] The pH of the composition is adjusted to a range of between about 3.0 and about 6.0, typically with an acid. Preferably, the pH is adjusted to a range of from about 5.0 to about 6.0. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

[0214] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0215] This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

[0216] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

[0217] This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene. As indicated above, other antioxidants such as butylated hydroxyanisole (BHA) can also be used.

[0218] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise one or more of methylparaben or propylparaben. Preferably, the preservative component comprises methylparaben and propylparaben. As indicated above, other preservatives can also be used.

[0219] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0220] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is adjusted to a range of from about 5.0 to about 6.0.

[0221] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 52.0% to about 80.0% of this embodiment of the composition. In an alternative in which the optimum concentration of allantoin is about 1.50% of the composition, the optimum concentration of water is about 61.65% of the composition. In another alternative, in which the optimum concentration of allantoin is about 9.00% of the composition, the optimum concentration of water is about 54.15% of the composition.

[0222] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0223] Sodium lauryl sulfate, as a 30% solution, can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, sodium lauryl sulfate, as a 30% solution, comprises from about 1.0% to about 3.0% of this embodiment of the composition. An optimum concentration of sodium lauryl sulfate, as a 30% solution, is about 1.90% of this embodiment of the composition.

[0224] Tetrasodium EDTA can comprise from about 0.05% to about 0.30% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

[0225] Galactoarabinan can comprise from about 1.0% to about 25.0% of this embodiment of the composition. Preferably, galactoarabinan comprises from about 3.0% to about 15.0% of this embodiment of the composition. An optimum concentration of galactoarabinan is about 5.00% of this embodiment of the composition.

[0226] Citric acid can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of citric acid is about 0.15% of this embodiment of the composition.

[0227] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0228] Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.20% of this embodiment of the composition.

[0229] Stearyl alcohol can comprise from about 0.50% to about 6.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 2.00% of this embodiment of the composition.

[0230] An acidic wax such as beeswax can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, the acidic wax such as beeswax comprises from about 1.0% to about 3.0% of this embodiment of the composition. An optimum concentration of the acidic wax such as beeswax is about 1.90% of this embodiment of the composition.

[0231] Cod liver oil can comprise from about 0.50% to about 15.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 10.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

[0232] Butylated hydroxytoluene can comprise from about 0.1% to about 3.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.25% to about 2.50% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

[0233] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0234] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0235] Allantoin can comprise from about 0.50% to about 10.0% of this embodiment of the composition. In one alternative, a preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. In this alternative, an optimum concentration of allantoin is about 1.50% of this embodiment of the composition. In another alternative, an optimum concentration of allantoin is about 9.00% of this embodiment of the composition.

[0236] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, if present, fragrance can comprise from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0237] Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin in a concentration of at least about 2.5%;

(2) an emulsifier system comprising:

- (a) an acidic wax; and
- (b) an anionic emulsifier that is substantially hydrophilic and is soluble in

water.

[0238] The acidic waxes used are as described above. A particularly preferred acidic wax is beeswax.

[0239] The anionic emulsifier that is substantially hydrophilic and soluble in water can be selected from the group consisting of ammonium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, ammonium lauryl sulfosuccinate, sodium dodecylbenzenesulfonate, ammonium laureth sulfate, sodium N-lauryl sarcosinate, and sodium lauryl sulfate. A particularly preferred anionic emulsifier is sodium lauryl sulfate.

[0240] The pH of the composition is adjusted to a range of between about 3.0 and about 6.0, typically with an acid. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

[0241] This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of ethylene glycol, propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

[0242] This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

[0243] This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, and stearyl alcohol. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and stearyl alcohol.

[0244] This embodiment of the invention can further comprise a preservative component. The preservative component can comprise one or more of methylparaben or propylparaben. Preferably, the preservative component comprises methylparaben and propylparaben. As indicated above, other preservatives can also be used.

[0245] This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

[0246] The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is adjusted to a range of from about 5.0 to about 6.0.

[0247] Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. An optimum concentration of water is about 58.98% of this embodiment of the composition

[0248] Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

[0249] Sodium lauryl sulfate, as a 30% solution, can comprise from about 0.50% to about 5.0% of this embodiment of the composition. An optimum concentration of sodium lauryl sulfate, as a 30% solution, is about 3.00% of this embodiment of the composition.

[0250] Tetrasodium EDTA can comprise from about 0.05% to about 0.50% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

[0251] Citric acid can comprise from about 0.05% to about 0.50% of this embodiment of the composition. An optimum concentration of citric acid is about 0.12% of this embodiment of the composition.

[0252] Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

[0253] Cetyl alcohol can comprise from about 3.0% to about 10.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.20% of this embodiment of the composition.

[0254] Stearyl alcohol can comprise from about 1.0% to about 5.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 2.00% of this embodiment of the composition.

[0255] An acidic wax such as beeswax can comprise from about 0.50% to about 5.0% of this embodiment of the composition. An optimum concentration of the acidic wax such as beeswax is about 3.00% of this embodiment of the composition.

[0256] Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

[0257] Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

[0258] Allantoin can comprise from about 2.5% to about 10.0% of this embodiment of the composition. An optimum concentration of allantoin is about 9.00% of this embodiment of the composition.

[0259] If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

[0260] Examples of particularly preferred compositions according to the present invention are described below.

[0261] Compositions according to the present invention can contain other, optional ingredients. For example, compositions according to the present invention can contain lipid-soluble components such as, but not limited to, caprylic/capric triglycerides; steareth-2; steareth-21; polyglyceryl-3 beeswax; a branched-carboxylic acid ester of a branched-chain alcohol selected from the group consisting of isononyl isononanoate, isodecyl isononanoate, isooctyl isononanoate, isononyl isooctanoate, isodecyl isooctanoate, isooctyl isooctanoate, isononyl isodecanoate, isooctyl isodecanoate, and isodecyl isodecanoate; an acrylates/C₁₀-C₃₀ alkyl acrylates cross-polymer; methylgluceth-20; a glyceryl ester of a long-chain fatty acid selected from the group consisting of glyceryl monostearate, glyceryl monopalmitate, and glyceryl monoarachidate; hydrogenated

vegetable oil; squalane; C₁₂-C₁₅ alkyl benzoates; di-C₁₂-C₁₅ alkyl fumarate; cholesterol; lanolin alcohol; octyldodecanol; isostearic acid; a branched-chain neopentanoate selected from the group consisting of octyldodecyl neopentanoate, heptyldodecyl neopentanoate, nonyldodecyl neopentanoate, octylundecyl neopentanoate, heptylundecyl neopentanoate, nonylundecyl neopentanoate, octyltridecyl neopentanoate, heptyltridecyl neopentanoate, and nonyltridecyl neopentanoate; an arachidyl ester of a short-chain carboxylic acid selected from the group consisting of arachidyl propionate, arachidyl acetate, arachidyl butyrate, and arachidyl isobutyrate; a long-chain fatty acid ester of a medium-chain alcohol selected from the group consisting of octyl palmitate, octyl myristate, octyl stearate, heptyl palmitate, heptyl myristate, heptyl stearate, nonyl palmitate, nonyl myristate, and nonyl stearate; jojoba oil; a myristyl ester of a long-chain fatty acid selected from the group consisting of myristyl myristate, myristyl laurate, and myristyl palmitate; bisabolol; hydrogenated jojoba oil; jojoba esters; methylgluceth-20 sesquisteate; PPG-14 butyl ether; PPG-15 stearyl ether; PPG-1-isoceteth-3-acetate; laureth-2-benzoate; diisostearyl dimer dilinoleate; a long-chain *cis*-monounsaturated fatty acid ester of a medium-chain alcohol; a medium-chain saturated carboxylic acid ester of a long-chain alcohol; hydrogenated soy glycerides; a long-chain fatty acid ester of cetyl alcohol selected from the group consisting of cetyl palmitate, cetyl stearate, and cetyl myristate; palm kernel oil; palm oil; and an arachidyl ester selected from the group consisting of arachidyl acetate, arachidyl propionate, arachidyl butyrate, and arachidyl isobutyrate.

[0262] In addition, the composition can further comprise other ingredients that are generally used in the cosmetic art and in the art of over-the-counter skin preparations. These ingredients include, but are not limited to: (1) other plant extracts, such as horsetail extract, horse chestnut extract, rose extract, or lavender extract; (2) a short-chain carboxylic acid ester of tocopherol selected from the group consisting of tocopheryl acetate, tocopheryl propionate, tocopheryl butyrate, and tocopheryl isobutyrate; (3) a long-chain fatty acid ester of ascorbic acid selected from the group consisting of ascorbyl myristate, ascorbyl palmitate, and ascorbyl stearate; (4) a long-chain fatty acid ester of retinol or a retinol derivative or analogue wherein the acyl moiety of the ester is selected from the group consisting of myristic acid, palmitic acid, and stearic acid; and (5), a sunscreen, which can be at least one compound selected from the group consisting of octyl

methoxycinnamate, *p*-aminobenzoic acid, ethyl *p*-aminobenzoate, isobutyl *p*-aminobenzoate, glyceryl *p*-aminobenzoate, *p*-dimethylaminobenzoic acid, methyl anthranilate, menthyl anthranilate, phenyl anthranilate, benzyl anthranilate, phenylethyl anthranilate, linalyl anthranilate, terpinyl anthranilate, cyclohexenyl anthranilate, amyl salicylate, phenyl salicylate, benzyl salicylate, menthyl salicylate, glyceryl salicylate, dipropyleneglycol salicylate, methyl cinnamate, benzyl cinnamate, α -phenyl cinnamionitrile, butyl cinnamoylpyruvate, umbelliferone, methylacetoumbelliferone, esculetin, methylesculetin, daphnetin, esculin, daphnin, diphenylbutadiene, stilbene, dibenzalacetone, benzalacetophenone, sodium 2-naphthol-3,6-disulfonate, sodium 2-naphthol-6,8-disulfonate, dihydroxynaphthoic acid, salts of dihydroxynaphthoic acid, *o*-hydroxybiphenyldisulfonates, *p*-hydroxybiphenyldisulfonates, 7-hydroxycoumarin, 7-methylcoumarin, 3-phenylcoumarin, 2-acetyl-3-bromoindazole, phenylbenzoxazole, methyl naphthoxazole, arylbenzothiazoles, quinine bisulfate, quinine sulfate, quinine chloride, quinine oleate, quinine tannate, 8-hydroxyquinoline salts, 2-phenylquinoline, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric acid, vilouric acid, tannic acid, tannic acid hexaethylether, hydroquinone, oxybenzone, sulisobenzene, dioxybenzone, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-dihydroxy-4,4'-dimethoxybenzophenone, octabenzene, 4-isopropyl dibenzoylmethane, butylmethoxydibenzoylmethane, etocrylene, and 4-isopropyl dibenzoylmethane.

[0263] Other ingredients can also optionally be included, such as colorants, pigments, opacifiers, and the like.

[0264] The composition is prepared by standard mixing techniques, such as are conventional in the cosmetic art and in the art of over-the-counter drug formulation for blending lipid-soluble components and water-soluble components. These mixing techniques include both manual and mechanical mixing, and include homogenization mixing and sweep mixing. The mixing techniques to be used can be chosen by one of ordinary skill in the art based on variables such as the viscosity of the components to be mixed and the volume of those components, as well as the relative proportion of lipid-soluble and water-soluble ingredients. The composition can be mixed in two or more batches, such as one batch containing lipid-soluble ingredients and another batch

containing water-soluble ingredients, and the batches can then be mixed at the final stage of preparation.

[0265] The batches to be used are shown in Tables 1-20.

[0266] Compositions according to the present invention can be formulated for the treatment of skin diseases and conditions. Among the skin diseases and conditions for which compositions according to the present invention can be formulated are epidermolysis bullosa, decubitus ulcers, pressure ulcers, diabetic ulcers, and milia. Compositions according to the present invention can be formulated for treatment of other skin diseases and conditions. The details of the composition can be varied according to the particular condition to be treated. For example, greater or lesser degrees of oil or lipid-soluble components can be included, and, in the case of compositions intended to be used on patients who may undergo allergic reactions, compounds that often generate allergic reactions, such as fragrance and coloring, can be excluded. Such details can readily be ascertained by one of ordinary skill in the art.

[0267] The invention is illustrated by the following Examples. These Examples are for illustrative purposes only and are not intended to limit the invention.

Example 1

Preparation of Skin Protectant Over-the-Counter Cream with pH of 7.4

(Prior Art Example)

[0268] A skin protectant over-the-counter (OTC) cream was prepared in accordance with the formulation of Table 1.

TABLE 1

COMPOSITION OF ALLANTOIN-CONTAINING SKIN CREAM WITH pH OF 7.4

<u>INGREDIENT</u>			<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
Part A					
Water			50.0-90.0	55.0-75.0	66.20
Sodium	Lauryl	Sulfate	0.50-2.50	1.00-2.50	1.90
(30%)					

Propylene Glycol	2.0-9.0	3.0-6.0	5.30
Tetrasodium EDTA	0.05-0.50	0.10-0.30	0.15

Part B

Lanolin Oil	5.0-15.0	8.0-12.0	10.60
Cetyl Alcohol	3.0-10.0	3.5-7.5	6.80
Stearyl Alcohol	1.0-5.0	1.0-3.0	2.00
Beeswax	0.50-2.50	1.0-2.5	1.90
Cod Liver Oil	1.0-7.0	1.0-4.0	2.00
BHT	0.10-1.00	0.20-0.80	0.50

Part C

St. John's Wort Extract	0.05-0.50	0.05-0.15	0.10
Witch Hazel Extract	0.05-0.50	0.05-0.15	0.10
Chamomile Extract	0.05-0.50	0.05-0.15	0.10
Arnica Extract	0.05-0.50	0.05-0.15	0.10
Methylparaben	0.10-0.50	0.15-0.40	0.30
Propylparaben	0.10-0.50	0.10-0.30	0.25
Allantoin	0.50-2.00	0.50-2.00	1.50
Fragrance	0.05-0.50	0.10-0.30	0.20

[0269] The Part A ingredients were combined and heated to 175°F with mixing. The Part B ingredients were combined and heated to 175°F with mixing. The Part B mixture was then added to the Part A mixture with mixing. The resulting mixture was then cooled to 120°F with continued mixing. The Part C ingredients were then added with mixing. The final emulsion was allowed to cool with continued mixing. The resulting cream had a pH of 7.4. Samples of the cream prepared from Example 1 were used for accelerated aging stability studies and analyzed for their allantoin concentration after a period of time at 40°C. The results are shown in Table 2.

[0270] As can be seen from Table 2, the allantoin in the cream from Example 1 undergoes degradation and would not meet the specifications required for an OTC drug.

TABLE 2
STABILITY OF ALLANTOIN IN SKIN CREAM COMPOSITION
OF EXAMPLE 1 WITH STORAGE AT 40°C

Days at 40°C	Weight % Allantoin
--------------	--------------------

0	1.5
30	1.4
60	1.3
90	1.2

CONFIDENTIAL

Example 2Preparation of a Cream Containing Allantoin with Lower pH

[0271] An OTC skin cream containing allantoin was prepared using the ingredients in Table 3 to provide a cream with a lower pH.

TABLE 3

**COMPOSITION OF ALLANTOIN-CONTAINING
SKIN CREAM WITH pH OF 5.3**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
Part A			
Water	50.0-90.0	55.0-75.0	68.68
Sodium Lauryl Sulfate (30%)	0.50-2.50	1.00-2.50	1.90
Propylene Glycol	2.0-9.0	3.0-6.0	5.30
Tetrasodium EDTA	0.05-0.50	0.10-0.30	0.15
Citric Acid	0.05-0.50	0.08-0.35	0.12
Part B			
Lanolin Oil	5.0-15.0	8.0-12.0	10.60
Cetyl Alcohol	3.0-10.0	3.5-7.5	4.20
Stearyl Alcohol	1.0-5.0	1.0-3.0	2.00
Beeswax	0.50-2.50	1.0-2.5	1.90
Cod Liver Oil	1.0-7.0	1.0-4.0	2.00
BHT	0.10-1.00	0.20-0.80	0.50
Part C			
St. John's Wort Extract	0.05-0.50	0.05-0.15	0.10
Witch Hazel Extract	0.05-0.50	0.05-0.15	0.10
Chamomile Extract	0.05-0.50	0.05-0.15	0.10
Arnica Extract	0.05-0.50	0.05-0.15	0.10
Methylparaben	0.10-0.50	0.15-0.40	0.30
Propylparaben	0.10-0.50	0.10-0.30	0.25
Allantoin	0.50-10.0	0.50-2.00	1.50
Fragrance	0.05-0.50	0.10-0.30	0.20

[0272] The Part A ingredients were combined and heated to 175°F with mixing. The Part B ingredients were combined and heated to 175°F with mixing. The Part B mixture was added to the Part A mixture with mixing. The resulting mixture was then cooled to 120°F with mixing at which time the Part C ingredients were added with mixing. The final emulsion was allowed to cool with continue mixing. The resulting cream had a pH of 5.3.

[0273] The citric acid is added to adjust the pH. Another acidic wax can be used to replace the beeswax as described above.

[0274] It was found that a similar cream was produced if Part B was added to Part A or Part A was added to Part B. However, the cream has a better appearance if the oil phase and water phase are homogenized under high shear after the two phases are added to one another.

[0275] Samples of the cream of this example were used for accelerated aging stability studies and analyzed for their allantoin concentration. The results are shown in Table 4. As can be seen from Table 4, the allantoin is stable over time in a cream with a pH of 5.3.

TABLE 4

**STABILITY OF ALLANTOIN IN SKIN CREAM COMPOSITION OF
EXAMPLE 2 WITH STORAGE AT 40°C**

Days at 40°C	Weight % Allantoin
0	1.4
30	1.4
60	1.4
90	1.4

Example 3

Preparation of Allantoin-Containing Skin Cream with Ionic Emulsifiers

[0276] An allantoin-containing skin cream with ionic emulsifiers is prepared according to Table 5. The preparation follows the method used in Example 2, with the ingredients in

each of Part A, Part B, and Part C being combined separately and then Part B being added to Part A, with Part C then being added to the combination of Part A and Part B. The pH is adjusted to a value in a range of from about 5.0 to about 5.8 by neutralizing the stearic acid with enough triethanolamine to reach this pH. Other bases can be used instead of triethanolamine.

TABLE 5**ALLANTOIN-CONTAINING SKIN CREAM WITH IONIC EMULSIFIERS**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 85.0	71.70
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Triethanolamine (99%)	0.20 - 4.0	0.50 - 3.0	1.25
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 7.0	2.0 - 6.0	3.50
Stearic Acid	0.50 - 5.0	1.0 - 4.0	2.50
Cod Liver Oil	1.0 - 7.0	1.5 - 5.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 4**Preparation of Allantoin-Containing Skin Cream with Lactylate Emulsifiers**

[0277] An allantoin-containing skin cream with the emulsifiers sodium stearoyl lactylate and sodium isostearoyl lactylate is prepared according to Table 6. The preparation follows the method used in Example 3. The pH is adjusted by the addition of the appropriate quantity of citric acid.

TABLE 6**ALLANTOIN-CONTAINING SKIN CREAM WITH LACTYLATE EMULSIFIERS**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 – 90.0	60.0 - 80.0	73.42
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Citric Acid	0.05 - 0.50	0.10 - 0.40	0.18
Sodium Stearoyl Lactylate	0.30 – 3.0	0.50 - 2.50	1.00
Sodium Isostearoyl Lactylate	0.05 – 1.0	0.10 - 0.70	0.25
Tetrasodium EDTA	0.05 - 0.25	0.10 - 0.20	0.15
<u>Part B</u>			
Lanolin Oil	5.0 – 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	3.80
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 – 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 – 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 5**Preparation of Allantoin-Containing Skin Cream with Carboxypolymethylene Polymer**

[0278] An allantoin-containing skin cream with carboxypolymethylene polymer is prepared according to Table 7. The preparation follows the method used in Example 3, except that the triethanolamine (Part D) is added last, after the combining of Parts A, B, and C, to avoid thickening of the emulsion. The triethanolamine is added to adjust the pH.

TABLE 7**ALLANTOIN-CONTAINING SKIN CREAM WITH
CARBOXYPOLYMETHYLENE POLYMER**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 – 90.0	60.0 - 80.0	73.55
Carboxypolymethylene Polymer	0.40 - 3.0	0.50 - 2.0	1.00
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.00
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	3.00
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 – 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20
<u>Part D</u>			
Triethanolamine (99%)	0.05 – 3.0	0.20 - 2.0	0.80

Example 6**Preparation of Allantoin-Containing Skin Cream with
Polyethylene Glycol Ethers of Cetearyl Alcohol**

[0279] An allantoin-containing skin cream with polyethylene glycol ethers of cetearyl alcohol is prepared according to Table 8. The preparation follows the method used in Example 3. The citric acid is added to adjust the pH.

TABLE 8

**ALLANTOIN-CONTAINING SKIN CREAM WITH
POLYETHYLENE GLYCOL ETHERS OF CETEARYL ALCOHOL**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 – 90.0	55.0 - 75.0	66.33
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Tetrasodium EDTA	0.05 - 0.50	0.10 - 0.30	0.15
Cetareth-25	0.50 – 4.0	2.00 - 3.50	2.60
Citric Acid	0.04 - 0.40	0.10 - 0.30	0.12
<u>Part B</u>			
Lanolin Oil	5.0 – 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	3.0 – 10.0	3.5 - 7.5	4.30
Stearyl Alcohol	1.0 - 5.0	2.0 - 4.0	3.50
Cetareth-6	0.50 – 4.0	1.0 - 3.0	1.80
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 – 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.50	0.10 - 0.30	0.15
Allantoin	0.50 – 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.30	0.20

Example 7

**Preparation of Allantoin-Containing Skin Cream with
Polyethylene Glycol Ester of Stearic Acid and Glyceryl Stearate**

[0280] An allantoin-containing skin cream with a polyethylene glycol ester of stearic acid and glyceryl stearate is prepared according to Table 9. The preparation follows the method used in Example 3. The citric acid is added to adjust the pH.

TABLE 9

**ALLANTOIN-CONTAINING SKIN CREAM WITH
POLYETHYLENE GLYCOL ESTER OF STEARIC ACID
AND GLYCERYL STEARATE**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 – 90.0	55.0 - 80.0	67.86
Propylene Glycol	2.0 - 9.0	4.3 - 7.0	5.70
Tetrasodium EDTA	0.05 - 0.50	0.10 - 0.30	0.15
Citric Acid	0.04 - 0.40	0.10 - 0.30	0.14
PEG-100 Stearate	1.0 - 5.0	1.5 - 3.0	2.60
<u>Part B</u>			
Lanolin Oil	5.0 – 15.0	2.0 - 12.0	10.60
Cetyl Alcohol	3.0 – 10.0	2.5 - 7.5	3.0
Stearyl Alcohol	1.0 - 4.0	1.0 - 3.5	2.50
Glyceryl Stearate	1.0 - 5.0	2.0 - 4.0	2.50
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 – 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.50	0.10 - 0.30	0.20
Allantoin	0.50 - 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 8

**Preparation of Allantoin-Containing Skin Cream with
Carboxypolymethylene Polymer and Polyethylene Glycol Ester of Stearic Acid**

[0281] An allantoin-containing skin cream with a carboxypolymethylene polymer and a polyethylene glycol ester of stearic acid is prepared according to Table 10. The preparation follows the method used in Example 5, with the triethanolamine (Part D) being added last. The triethanolamine is added to adjust the pH.

TABLE 10

**ALLANTOIN-CONTAINING SKIN CREAM WITH
A CARBOXYPOLYMETHYLENE POLYMER AND A
POLYETHYLENE GLYCOL ESTER OF STEARIC ACID**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 85.0	69.95
Carboxypolymethylene	0.30 - 3.0	0.50 - 2.0	0.85
Polymer	2.0 - 9.0	4.0 - 7.0	5.70
Propylene Glycol	0.25 - 2.5	0.50 - 2.0	1.50
PEG-100 Stearate			
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	4.20
Stearyl Alcohol	0.50 - 6.0	0.75 - 5.0	1.50
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.25	0.10 - 0.20	0.15
Allantoin	0.50 - 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20
<u>Part D</u>			
Triethanolamine (99%)	0.05 - 3.0	0.20 - 2.0	0.80

Example 9

**Preparation of Allantoin-Containing Skin Cream with
Galactoarabinan, Sodium Lauryl Sulfate, and Beeswax**

[0282] An allantoin-containing skin cream with galactoarabinan, sodium lauryl sulfate, and beeswax is prepared according to Table 11. The preparation follows the method used in Example 3. The citric acid is used to adjust the pH. Another acidic wax can substitute for beeswax.

TABLE 11

**ALLANTOIN-CONTAINING SKIN CREAM WITH
GALACTOARABINAN, SODIUM LAURYL SULFATE,
AND BEESWAX**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 80.0	61.65
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Sodium Lauryl Sulfate (30%)	0.50 - 5.0	1.0 - 3.0	1.90
Tetrasodium EDTA	0.05 - 0.30	0.10 - 0.20	0.15
Galactoarabinan	1.0 - 25.0	3.0 - 15.0	5.00
Citric Acid	0.05 - 0.25	0.10 - 0.20	0.15
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	4.20
Stearyl Alcohol	0.50 - 6.0	1.0 - 4.0	2.00
Beeswax	0.50 - 5.0	1.0 - 3.0	1.90
Cod Liver Oil	0.50 - 15.0	1.0 - 10.0	2.00
Butylated Hydroxytoluene	0.10 - 3.0	0.25 - 2.5	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 10.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 10

Preparation of a Cream Containing Allantoin with pH of 5.3 with High
Allantoin Concentration

[0283] An OTC skin cream containing allantoin is prepared using the ingredients in Table 12 to provide a cream with a lower pH with an allantoin concentration of about 9.00%. The skin cream is prepared according to the method of Example 2.

TABLE 12

**COMPOSITION OF ALLANTOIN-CONTAINING SKIN
CREAM WITH pH OF 5.3 WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	61.38
Sodium Lauryl Sulfate (30%)	0.50-2.50	1.90
Propylene Glycol	2.0-9.0	5.30
Tetrasodium EDTA	0.05-0.50	0.15
Citric Acid	0.05-0.50	0.12
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	3.0-10.0	4.20
Stearyl Alcohol	1.0-5.0	2.00
Beeswax	0.50-2.50	1.90
Cod Liver Oil	1.0-7.0	2.00
BHT	0.10-1.00	0.50
Part C		
St. John's Wort Extract	0.05-0.50	0.10
Witch Hazel Extract	0.05-0.50	0.10
Chamomile Extract	0.05-0.50	0.10
Arnica Extract	0.05-0.50	0.10
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Allantoin	0.50-10.00	9.00
Fragrance	0.05-0.50	0.20

Example 11

**Preparation of Allantoin-Containing Skin Cream with Ionic Emulsifiers with High
Allantoin Concentration**

[0284] An OTC skin cream containing allantoin is prepared using the ingredients in Table 13 to provide a cream with an allantoin concentration of about 9.00% using ionic emulsifiers. The skin cream is prepared according to the method of Example 3. The pH is

adjusted to a value in a range of from about 5.0 to about 5.8 by neutralizing the stearic acid with enough triethanolamine to reach this pH. Other bases can be used instead of triethanolamine.

CONFIDENTIAL

TABLE 13

ALLANTOIN-CONTAINING SKIN CREAM WITH IONIC EMULSIFIERS
WITH HIGH ALLANTOIN CONCENTRATION

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	64.20
Propylene Glycol	2.0-9.0	5.70
Triethanolamine (99%)	0.20-4.0	1.25
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	1.0-7.0	3.50
Stearic Acid	0.50-5.0	2.50
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Allantoin	0.50-10.0	9.00
Fragrance	0.05-0.50	0.20

Example 12

Preparation of a Cream Containing Allantoin with Lactylate Emulsifiers with High
Allantoin Concentration

[0285] An OTC skin cream containing allantoin is prepared using the ingredients in Table 14 to provide a cream with an allantoin concentration of about 9.00% using lactylate emulsifiers. The skin cream is prepared according to the method of Example 4. The pH is adjusted by the addition of the appropriate quantity of citric acid.

TABLE 14**ALLANTOIN-CONTAINING SKIN CREAM WITH LACTYLATE
EMULSIFIERS WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	65.92
Propylene Glycol	2.0-9.0	5.70
Citric Acid	0.05-0.50	0.18
Sodium Stearoyl Lactylate	0.30-3.0	1.00
Sodium Isostearoyl Lactylate	0.05-1.0	0.25
Tetrasodium EDTA	0.05-0.25	0.15
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	1.0-8.0	3.80
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Allantoin	0.50-1.00	9.00
Fragrance	0.05-0.50	0.20

Example 13**Preparation of a Cream Containing Allantoin with Carboxypolymethylene
Polymer with High Allantoin Concentration**

[0286] An OTC skin cream containing allantoin is prepared using the ingredients in Table 15 to provide a cream with an allantoin concentration of about 9.00% with a carboxypolymethylene polymer. The skin cream is prepared according to the method of Example 5. Triethanolamine is added to adjust the pH.

TABLE 15**ALLANTOIN-CONTAINING SKIN CREAM WITH CARBOXYPOLYMETHYLENE
POLYMER WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	66.05
Carboxypolymethylene Polymer	0.40-3.0	1.00
Propylene Glycol	2.0-9.0	5.70
Part B		
Lanolin Oil	5.0-15.0	10.00
Cetyl Alcohol	1.0-8.0	3.00
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Allantoin	0.50-10.0	9.00
Fragrance	0.05-0.50	0.20
Part D		
Triethanolamine (99%)	0.05-3.0	0.80

Example 14**Preparation of a Cream Containing Allantoin with Polyethylene Glycol Ethers of Cetearyl
Alcohol with High Allantoin Concentration**

[0287] An OTC skin cream containing allantoin is prepared using the ingredients in Table 16 to provide a cream with an allantoin concentration of about 9.00% with polyethylene glycol ethers of cetearyl alcohol. The skin cream is prepared according to the method of Example 6. The citric acid is added to adjust the pH.

TABLE 16

**ALLANTOIN-CONTAINING SKIN CREAM WITH POLYETHYLENE
GLYCOL ETHERS OF CETEARYL ALCOHOL WITH
HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	58.83
Propylene Glycol	2.0-9.0	5.70
Tetrasodium EDTA	0.05-0.50	0.15
Ceteareth-25	0.50-4.0	2.60
Citric Acid	0.04-0.40	0.12
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	3.0-10.0	4.30
Stearyl Alcohol	1.0-5.0	3.50
Ceteareth-6	0.50-4.0	1.80
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Diazolidinyl Urea	0.05-0.50	0.15
Allantoin	0.50-10.0	9.00
Fragrance	0.05-0.50	0.20

Example 15

**Preparation of a Cream Containing Allantoin with Polyethylene Glycol Ethers of Stearic
Acid and Glyceryl Stearate with High Allantoin Concentration**

[0288] An OTC skin cream containing allantoin is prepared using the ingredients in Table 17 to provide a cream with an allantoin concentration of about 9.00% with polyethylene glycol ethers of stearic acid and glyceryl stearate. The skin cream is prepared according to the method of Example 7. The citric acid is added to adjust the pH.

TABLE 17

**ALLANTOIN-CONTAINING SKIN CREAM WITH POLYETHYLENE
GLYCOL ESTER OF STEARIC ACID AND GLYCERYL STEARATE
WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	60.36
Propylene Glycol	2.0-9.0	5.70
Tetrasodium EDTA	0.05-0.50	0.15
Citric Acid	0.04-0.40	0.14
PEG-100 Stearate	1.0-5.0	2.60
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	3.0-10.0	4.30
Stearyl Alcohol	1.0-5.0	3.50
Glyceryl Stearate	1.0-5.0	2.50
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Diazolidinyl Urea	0.05-0.50	0.20
Allantoin	0.50-10.0	9.00
Fragrance	0.05-0.50	0.20

Example 16

**Preparation of a Cream Containing Allantoin with a Carboxypolymethylene Polymer and
a Polyethylene Glycol Ether of Stearic Acid with High Allantoin Concentration**

[0289] An OTC skin cream containing allantoin is prepared using the ingredients in Table 18 to provide a cream with an allantoin concentration of about 9.00% with a carboxypolymethylene polymer and a polyethylene glycol ester of stearic acid. The skin cream is prepared according to the method of Example 8. The triethanolamine is added to adjust the pH.

TABLE 18

**ALLANTOIN-CONTAINING SKIN CREAM WITH A CARBOXYPOLYMETHYLENE
POLYMER AND A POLYETHYLENE GLYCOL ESTER OF STEARIC ACID
WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	62.45
Carboxypolymethylene Polymer	0.30-3.0	0.85
Propylene Glycol	2.0-9.0	5.70
PEG-100 Stearate	0.25-2.5	1.50
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	1.0-8.0	4.20
Stearyl Alcohol	0.50-6.0	1.50
Cod Liver Oil	1.0-7.0	2.00
Butylated Hydroxytoluene	0.10-1.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Diazolidinyl Urea	0.05-0.25	0.15
Allantoin	0.50-9.0	9.00
Fragrance	0.05-0.50	0.20
Part D		
Triethanolamine (99%)	0.05-3.0	0.80

Example 17

Preparation of a Cream Containing Allantoin with Galactoarabinan, Sodium Lauryl Sulfate,
and Beeswax with High Allantoin Concentration

[0290] An OTC skin cream containing allantoin is prepared using the ingredients in Table 19 to provide a cream with an allantoin concentration of about 9.00% with galactoarabinan, sodium lauryl sulfate, and beeswax. The skin cream is prepared according to the method of Example 9. The citric acid is used to adjust the pH. Another acidic wax can substitute for beeswax.

TABLE 19

**COMPOSITION OF ALLANTOIN-CONTAINING SKIN
CREAM WITH GALACTOARABINAN, SODIUM LAURYL SULFATE,
AND BEESWAX WITH HIGH ALLANTOIN CONCENTRATION**

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	54.15
Propylene Glycol	2.0-9.0	5.70
Sodium Lauryl Sulfate (30%)	0.50-5.0	1.90
Tetrasodium EDTA	0.05-0.30	0.15
Galactoarabinan	1.0-25.0	5.00
Citric Acid	0.05-0.25	0.15
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	1.0-8.0	4.20
Stearyl Alcohol	0.50-6.0	2.00
Beeswax	0.50-5.0	1.90
Cod Liver Oil	0.50-15.0	2.00
Butylated Hydroxytoluene	0.10-3.0	0.50
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Allantoin	0.50-10.0	9.00
Fragrance	0.05-0.50	0.20

Example 18

**Preparation of a Cream Containing Allantoin with Sodium Lauryl Sulfate,
and Beeswax with High Allantoin Concentration and pH of 3.9**

[0291] An OTC skin cream containing allantoin is prepared using the ingredients in Table 20 to provide a cream with an allantoin concentration of about 9.00% with a pH of 3.9 with sodium lauryl sulfate and beeswax. The water phase (Part A in Table 20) was heated to 160-180°F. The oil phase (Part B in Table 20) was heated to 160-180°F. The heated oil phase was added to the heated water phase with continuing mixing to form an oil-in-water

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

TABLE 20

COMPOSITION OF ALLANTOIN-CONTAINING SKIN CREAM
WITH pH OF 3.9 WITH BEESWAX AND HIGH ALLANTOIN CONCENTRATION

<u>INGREDIENT</u>	<u>RANGE</u>	<u>OPTIMUM</u>
Part A		
Water	50.0-90.0	58.98
Sodium Lauryl Sulfate (30%)	0.50-5.0	3.00
Propylene Glycol	2.0-9.0	5.70
Tetrasodium EDTA	0.05-0.50	0.15
Citric Acid	0.05-0.50	0.12
Part B		
Lanolin Oil	5.0-15.0	10.60
Cetyl Alcohol	3.0-10.0	4.20
Stearyl Alcohol	1.0-5.0	2.00
Beeswax	0.50-5.0	3.00
Part C		
Methylparaben	0.10-0.50	0.30
Propylparaben	0.10-0.50	0.25
Fragrance	0.05-0.50	0.20
Allantoin	2.50-10.0	9.00

ADVANTAGES OF THE PRESENT INVENTION

[0292] The present invention provides an allantoin-containing composition that is an oil-in-water emulsion using either an acidic anionic polymer and an anionic emulsifier or an acidic anionic polymer and a nonionic emulsifier that is an ethoxylated ether or ethoxylated ester. The composition can further include a carbohydrate polymer that can be polygalactose or polyarabinose. If an ethoxylated ether or ethoxylated ester is used, the composition can further comprise glyceryl stearate. The composition has improved thermal stability. The composition according to the present invention is useful for treatment of a number of skin diseases and conditions, including epidermolysis bullosa, pressure ulcers, diabetic ulcers, decubitus ulcers, and milia, as well as other inflammatory conditions, such as conditions

[0293] Although the present invention has been described in considerable detail, with reference to certain preferred versions thereof, other versions and embodiments are possible. Therefore, the scope of the invention is determined by the following claims.

-65-